Brain Research, 493 (1989) 407–409
Elsevier

Author Index

Amrbein, C.L., see Surichamorn, W., 320
Aoki, C., Zemcik, B.A., Strader, C.D. and Pickel, V.M., Cytoplasmic loop of β-adrenergic receptors: synaptic and intracellular localization and relation to catecholaminergic neurons in the nuclei of the solitary tracts, 331
Ashby Jr., C.R., Hitzemann, R., Rubinstein, J.E. and Wang, R.Y., One year treatment with haloperidol or clozapine fails to alter neostriatal D_1- and D_2-dopamine receptor sensitivity in the rat, 194

Baimbridge, K.G., see Mudrick, L.A., 179
Balbo, A., see Wadhwani, K.C., 251
Bandler, R., see Carrive, P., 385
Barnes, M.I., see Rogers, B.C., 41
Bauswein, E., Fromm, C. and Preuss, A., Corticostriatal cells in comparison with pyramidal tract neurons: contrasting properties in the behaving monkey, 198
Bidlack, J.M., see Selley, D.E., 23
Bland, B.H., see Ford, R.D., 269
Bledsoe Jr., S.C., McLaren, J.D. and Meyer, J.R., Potassium-induced release of endogenous glutamate and two as yet unidentified substances from the lateral line of *Xenopus laevis*, 113
Bolam, J.P., see Smith, Y., 160
Bouras, C., see Hof, P.R., 368
Bramham, C.R. and Srebro, D., Synaptic plasticity in the hippocampus is modulated by behavioral state, 74
Bullitt, E., Induction of c-*fos*-like protein within the lumbar spinal cord and thalamus of the rat following peripheral stimulation, 391
Buzsáki, G., see Masliah, E., 240

Carrive, P., Bandler, R. and Dampney, R.A.L., Viscerotopic control of regional vascular beds by discrete groups of neurons within the midbrain periaqueductal gray, 385
Castro, A.J., see Neafsey, E.J., 33
Chang, J.W., see Cornwell-Jones, C.A., 258
Chavkin, C., see Neumaier, J.F., 292
Cole, B., see Cornwell-Jones, C.A., 258
Colom, L.V., see Ford, R.D., 269
Confort-Gouny, S., see Vion-Dury, J., 175
Constantinidis, J., see Hof, P.R., 368
Cooper, A.J.L., see Raps, S.P., 398
Cornwell-Jones, C.A., Decker, M.W., Chang, J.W., Cole, B., Goltz, K.M., Tran, T. and McGaugh, J.L., Neonatal 6-hydroxydopa, but not DSP-4, elevates brainstem monoamines and impairs inhibitory avoidance learning in developing rats, 258
Coughlin, C. and Elam, J.S., Enhanced axonal transport of glycosaminoglycans in regenerating goldfish optic nerve, 326
Cozzone, P.J., see Vion-Dury, J., 175

Dampney, R.A.L., see Carrive, P., 385
Davidoff, R.A., see Zhang, D.X., 129
Decker, M.W., see Cornwell-Jones, C.A., 258
Denton, D.A., see Park, R., 123
Ducis, I., Norenberg, L.-O.B. and Norenberg, M.D., Effect of ammonium chloride on the astrocyte benzodiazepine receptor, 362
Durmus, G., see Öcal-İrez, T., 1

Elam, J.S., see Coughlin, C., 326
El-Fakahany, E.E., see Surichamorn, W., 320
Emadian, S.M., see Luttge, W.G., 190
Enna, S.J., see Pilc, A., 8

Fantini, J., see Vion-Dury, J., 175
Ford, R.D., Colom, L.V. and Bland, B.H., The classification of medial septum-diagonal band cells as θ-on or θ-off in relation to hippocampal EEG states, 269
Forray, C., see Surichamorn, W., 320
Fromm, C., see Bauswein, E., 198
Fukuda, A., Nabekura, J., Ito, C.R., Plata-Salamán, C.R. and Oomura, Y., Developmentally different onset of α_1- and α_2-adrenergic responses in the neonatal rat dorsal motor nucleus of the vagus in vitro, 357
Fukuda, M., see Miyake, T., 376
Futenma, A., see Gotoh, M., 97

Gale, K., see Segovia, J., 185
Galons, J.-P., see Vion-Dury, J., 175
Goltz, K.M., see Cornwell-Jones, C.A, 258
Gotoh, M., Iguchi, A., Yatomi, A., Uemura, K., Miura, H., Futenma, A., Kato, K. and Sakamoto, N., Vagally mediated insulin secretion by stimulation of brain cholinergic neurons with neostigmine in bilateral adrenalectomized rats, 97
Gulyàs, B., see Raiguel, S.E., 155

Hackman, J.C., see Zhang, D.X., 129
Harrington, M.E., see Rusak, B., 283
Harvey, J.A., see Winksy, L., 136
Hertz, L., see Raps, S.P., 398
Higashima, M., see Masukawa, L.M., 168
Hitzemann, R., see Ashby Jr., C.R., 194
Hof, P.R., Bouras, C., Constantinidis, J. and Morrison, J.H., Balint's syndrome in Alzheimer's disease: specific disruption of the occipito-parietal visual pathway, 368
Hughes, D., see Smith, S.M., 366

Iguchi, A., see Gotoh, M., 97
Imamura, Y., see Miyake, T., 376
Inoue, N., Magari, S., Ito, Y. and Sakanaka, M., Distribution, possible origins and fine structure of neuropeptide Y-containing nerve fibers in the rat liver, 87
Ishikawa, H., see Sakuma, S., 402
Ito, C.R., see Fukuda, A., 357
Ito, Y., see Inoue, N., 87
Iwahasi, K., see Ozaki, H.S., 66

Jacobowitz, D.M., see Winksy, L., 136
Jahnke, M.T., Proske, U. and Struppler, A., Measurements of muscle stiffness, the electromyogram and activity in single muscle spindles of human flexor muscles following conditioning by passive stretch or contraction, 103

Joyce, J.N., see Marshall, J.F., 247

Kang, C.G., see Luttge, W.G., 190
Kato, K., see Gotoh, M., 97
Kenigfest, N.B., see Vesselkin, N.P., 51
Kim, J.H., see Masukawa, L.M., 168
Kitamura, T., see Miyake, T., 376
Kriat, M., see Vion-Dury, J., 175

Lagae, L., see Raiguel, S.E., 155
Lai, J.C.K., see Raps, S.P., 398
Lakos, S.F., see Miller, A.D., 348
Latker, C.H., see Wadhwani, K.C., 231
Low, P.A., see Shupeck, M., 225
Luttge, W.G., Kang, C.G., Rupp, M.E. and Emadian, S.M., Treatment of mouse brain cytosol with dextran-coated charcoal and high salt does not reveal a new glucocorticoid binder, 190

Maekawa, K., see Sakuma, S., 402
Magari, S., see Inoue, N., 87
Marshall, J.F., Navarrete, R. and Joyce, J.N., Decreased striatal D_1 binding density following mesotelencephalic 6-hydroxydopamine injections: an autoradiographic analysis, 247
Masliah, E., Terry, R. and Buzsáki, G., Thalamic nuclei in Alzheimer disease: evidence against the cholinergic hypothesis of plaque formation, 240
Masukawa, L.M., Higashima, M., Kim, J.H. and Spencer, D.D., Epileptiform discharges evoked in hippocampal brain slices from epileptic patients, 68
McGaugh, J.L., see Cornwell-Jones, C.A., 258
McKinley, M.J., see Park, R., 123
McLaren, J.D., see Bledsoe Jr., S.C., 113
McMaster, S.E., see Winksy, L., 136
Meijer, J.H., see Rusak, B., 283
Meinecke, D.L., Tallman, J. and Rakic, P., $GABA_A$/benzodiazepine receptor-like immunoreactivity in rat and monkey cerebellum, 303
Meloni, R., see Segovia, J., 185
Meyer, J.R., see Bledsoe Jr., S.C., 113
Miceli, D., see Vesselkin, N.P., 51
Miller, A.D., Tan, L.K. and Lakos, S.F., Brainstem projections to cats' upper lumbar spinal cord: implications for abdominal muscle control, 348
Mitchell, C.L., see Rogers, B.C., 41
Miura, H., see Gotoh, M., 97
Miyake, T., Imamura, Y., Fukuda, M. and Kitamura, T., Delay of astrocyte reaction in the injured cerebral cortex of hypothyroid mouse, 376
Morrison, J.H., see Hof, P.R., 368
Mudrick, L.A. and Baimbridge, K.G., Long-term structural changes in the rat hippocampal formation following cerebral ischemia, 179

Nabekura, J., see Fukuda, A., 357
Navarrete, R., see Marshall, J.F., 247
Neafsey, E.J., Sørensen, J.C., Tønder, N. and Castro, A.J., Fetal cortical transplants into neonatal rats respond to thalamic and peripheral stimulation in the adult. An electrophysiological study of single-unit activity, 33
Neumaier, J.F. and Chavkin, C., Release of endogenous opioid peptides displaces [^{3}H]diprenorphine binding in rat hippocampal slices, 292
Nomura, S., see Pilc, A., 8
Norenberg, L.-O.B., see Ducis, I., 362
Norenberg, M.D., see Ducis, I., 362

Öcal-İrez, T., Durmus, G., Sekerkiran, Y., Peker, C. and Uygur, C., Effect of a pineal indolamine, 5-methoxyindole-3-acetic acid, on the estrous cycle and reproductive organs of female Wistar albino rats, 1
Ono, J.K., Synaptic connections in the buccal ganglia of *Aplysia* mediated by an identified neuron containing a CCK/gastrin-like peptide co-localized with acetylcholine, 212
Oomura, Y., see Fukuda, A., 357
Orban, G.A., see Raiguel, S.E., 155
Ozaki, H.S., Iwahashi, K. and Shimada, M., Ipsilateral corticocortical projections of fibers which course within Probst's longitudinal bundle seen in the brains of mice with congenital absence of the corpus callosum: a study with the horseradish peroxidase technique, 66

Park, R., Denton, D.A., McKinley, M.J., Pennington, G. and Weisinger, R.S., Intracerebroventricular saccharide infusions inhibit thirst induced by systemic hypertonicity, 123
Peker, C., see Öcal-İrez, T., 1
Pennington, G., see Park, R., 123
Pickel, V.M., see Aoki, C., 331
Pilc, A., Vetulani, J., Nomura, S. and Enna, S.J., An in vivo procedure for the selective inactivation of rat brain cerebral cortical α-adrenoceptors using *N*-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline (EEDQ), 8
Plata-Salamán, C.R., see Fukuda, A., 357
Polakiewicz, R.D., see Rosen, H., 380
Prado, W.A., Antinociceptive effect of agonists microinjected into the anterior pretectal nucleus of the rat, 147
Preuss, A., see Bauswein, E., 198
Proske, U., see Jahnke, M.T., 103

Raiguel, S.E., Lagae, L., Gulyàs, B. and Orban, G.A., Response latencies of visual cells in macaque areas V1, V2 and V5, 155
Rakic, P., see Meinecke, D.L., 303
Rapoport, S.I., see Wadhwani, K.C., 231
Raps, S.P., Lai, J.C.K., Hertz, L. and Cooper, A.J.L., Glutathione is present in high concentrations in cultured astrocytes but not in cultured neurons, 398
Repérant, J., see Vesselkin, N.P., 51
Rio, J.P., see Vesselkin, N.P., 51
Rogers, B.C., Barnes, M.I., Mitchell, C.L. and Tilson, H.A., Functional deficits after sustained stimulation of the perforant path, 41
Rosen, H. and Polakiewicz, R.D., Increase in hypothalamic pro-opiomelanocortin gene expression in response to prolonged low level lead exposure, 380
Rubinstein, J.E., see Ashby Jr., C.R., 194
Rupp, M.E., see Luttge, W.G., 190
Rusak, B., Meijer, J.H. and Harrington, M.E., Hamster circadian rhythms are phase-shifted by electrical stimulation of the geniculo-hypothalamic tract, 283

Sakamoto, N., see Gotoh, M., 97
Sakanaka, M., see Inoue, N., 87
Sakuma, S., Maekawa, K. and Ishikawa, H., Immunohistochemical studies on the somatostatin- and growth hormone-releasing factor (GRF)-neurons in the hypothalamus of the novel dwarf rat; the spontaneous dwarf rat (*dr*), 402

Scharfman, H.E. and Schwartzkroin, P.A., Selective depression of GABA-mediated IPSPs by somatostatin in area CA1 of rabbit hippocampal slices, 205
Schmelzer, J.D., see Shupeck, M., 225
Schwartzkroin, P.A., see Scharfman, H.E., 205
Segovia, J., Meloni, R. and Gale, K., Effect of dopaminergic denervation and transplant-derived reinnervation on a marker of striatal GABAergic function, 185
Sekerkiran, Y., see Öral-İrez, T., 1
Selley, D.E. and Bidlack, J.M., Guanine nucleotide regulation of [^{125}I]β-endorphin binding to NG108-15 and SK-N-SH cell membranes: specific cation requirements, 23
Shimada, M., see Ozaki, H.S., 66
Shupeck, M., Ward, K.K., Schmelzer, J.D. and Low, P.A., Comparison of nerve regeneration in vascularized and conventional grafts: nerve electrophysiology, norepinephrine, prostacyclin, malondialdehyde, and the blood–nerve barrier, 225
Shupliakov, O.V., see Vesselkin, N.P., 51
Smith, D.O., see Williams, K.D., 14
Smith, S.M. and Hughes, D., 1-(2-Pyrimidinyl)piperazine antagonizes GABA-activated currents in cultured spinal neurones of the rat, 366
Smith, Y. and Bolam, J.P., Neurons of the substantia nigra reticulata receive a dense GABA-containing input from the globus pallidus in the rat, 160
Sørensen, J.C., see Neafsey, E.J., 33
Spencer, D.D., see Masukawa, L.M., 168
Srebro, B., see Bramham, C.R., 74
Strader, C.D., see Aoki, C., 331
Struppler, A., see Jahnke, M.T., 103
Surichamorn, W., Amrhein, C.L., Forray, C. and El-Fakahany, E.E., Inhibition of cyclic AMP formation in N1E-115 neuroblastoma cells is mediated by a non-cardiac M2 muscarinic receptor subtype, 320

Tallman, J., see Meinecke, D.L., 303
Tan, L.K., see Miller, A.D., 348
Terry, R., see Masliah, E., 240
Tilson, H.A., see Rogers, B.C., 41
Tønder, N., see Neafsey, E.J., 33
Tran, T., see Cornwell-Jones, C.A., 258

Uemura, K., see Gotoh, M., 97
Uygur, G., see Öcal-İrez, T., 1

Vesselkin, N.P., Repérant, J., Kenigfest, N.B., Rio, J.P., Miceli, D. and Shupliakov, O.V., Centrifugal innervation of the lamprey retina. Light- and electron microscopic and electrophysiological investigations, 51
Vetulani, J., see Pilc, A., 8
Vion-Dury, J., Galons, J.-P., Fantini, J., Confort-Gouny, S., Kriat, M. and Cozzone, P.J., Phosphorus-31 nuclear magnetic resonance study of the C6 glioma cell line cultured on microcarrier beads, 175

Wadhwani, K.C., Latker, C.H., Balbo, A. and Rapoport, S.I., Perineurial permeability and endoneurial edema during Wallerian degeneration of the frog peripheral nerve, 231
Wang, R.Y., see Ashby Jr., C.R., 194
Ward, K.K., see Shupeck, M., 225
Weisinger, R.S., see Park, R., 123
Williams, K.D. and Smith, D.O., Cholesterol conservation in skeletal muscle associated with age- and denervation-related atrophy, 14
Winksy, L., Harvey, J.A., McMaster, S.E. and Jacobowitz, D.M., A study of proteins in the auditory system of rabbits using two-dimensional gels: identification of glial fibrillary acidic protein and vitamin D-dependent calcium binding protein, 136

Yatomi, A., see Gotoh, M., 97

Zemcik, B.A., see Aoki, C., 331
Zhang, D.X., Hackman, J.C. and Davidoff, R.A., NMDA antagonists and potentiation of NMDA-induced motoneuron depolarizations in the isolated frog spinal cord, 129